



Design and Analysis of Algorithms

Mohammad GANJTABESH

`mgtabesh@ut.ac.ir`

School of Mathematics, Statistics and Computer Science,
University of Tehran,
Tehran, Iran.


Solving the recurrence equations

There are different approaches to do this:

- Constructing Recursion Tree
- Performing Substitution
- Using Induction
- Master Theorem
- Generating Functions


Generating Functions

Generating Functions transform problems about sequences into problems about functions where we have powerful Mathematical tools.



Generating Functions


Generating Functions transform problems about sequences into problems about functions where we have powerful Mathematical tools.



$$\langle s_0, s_1, s_2, s_3, \dots \rangle \longleftrightarrow F(x) = \sum_{i=0}^{\infty} s_i x^i$$

Generating Functions

Generating Functions transform problems about sequences into problems about functions where we have powerful Mathematical tools.



$$\langle s_0, s_1, s_2, s_3, \dots \rangle \longleftrightarrow F(x) = \sum_{i=0}^{\infty} s_i x^i$$

Examples:

$$\langle 1, 1, 1, 1, \dots \rangle \longleftrightarrow 1 + z + z^2 + z^3 + \dots = \frac{1}{1 - z}$$

Generating Functions

Example

- $\langle 0, 0, 0, 0, \dots \rangle \longleftrightarrow 0 + 0x + 0x^2 + 0x^3 + \dots = 0.$
- $\langle 1, 0, 0, 0, \dots \rangle \longleftrightarrow 1 + 0x + 0x^2 + 0x^3 + \dots = 1.$
- $\langle 3, 2, 1, 0, \dots \rangle \longleftrightarrow 3 + 2x + x^2 + 0x^3 + \dots = 3 + 2x + x^2.$
- $\langle 1, -1, 1, -1, \dots \rangle \longleftrightarrow 1 - x + x^2 - x^3 + \dots = \frac{1}{1+x}.$
- $\langle 1, a, a^2, a^3, \dots \rangle \longleftrightarrow 1 + ax + a^2x^2 + a^3x^3 + \dots = \frac{1}{1-ax}.$
- $\langle 1, 0, 1, 0, \dots \rangle \longleftrightarrow 1 + x^2 + x^4 + \dots = \frac{1}{1-x^2}.$

Operations on Generating Functions

Scaling

if

$$\langle f_0, f_1, f_2, f_3, \dots \rangle \longleftrightarrow F(x),$$

then

$$\langle cf_0, cf_1, cf_2, cf_3, \dots \rangle \longleftrightarrow cF(x).$$

Operations on Generating Functions

Scaling

if

$$\langle f_0, f_1, f_2, f_3, \dots \rangle \longleftrightarrow F(x),$$

then

$$\langle cf_0, cf_1, cf_2, cf_3, \dots \rangle \longleftrightarrow cF(x).$$

Addition

if

$$\langle f_0, f_1, f_2, f_3, \dots \rangle \longleftrightarrow F(x)$$

and

$$\langle g_0, g_1, g_2, g_3, \dots \rangle \longleftrightarrow G(x),$$

then

$$\langle f_0 + g_0, f_1 + g_1, f_2 + g_2, f_3 + g_3, \dots \rangle \longleftrightarrow F(x) + G(x).$$

Operations on Generating Functions

Example

$$\langle 1, 0, 1, 0, \dots \rangle \longleftrightarrow 1 + x^2 + x^4 + \dots = \frac{1}{1 - x^2}$$

$$\implies \frac{2}{1 - x^2} = 2 + 2x^2 + 2x^4 + \dots$$

Operations on Generating Functions

Example

$$\begin{aligned}\langle 1, 0, 1, 0, \dots \rangle &\longleftrightarrow 1 + x^2 + x^4 + \dots = \frac{1}{1 - x^2} \\ \implies \frac{2}{1 - x^2} &= 2 + 2x^2 + 2x^4 + \dots\end{aligned}$$

Example

$$\langle 1, 1, 1, 1, \dots \rangle \longleftrightarrow \frac{1}{1 - x}$$

and

$$\langle 1, -1, 1, -1, \dots \rangle \longleftrightarrow \frac{1}{1 + x}$$

$$\implies \langle 2, 0, 2, 0, \dots \rangle \longleftrightarrow \frac{1}{1 - x} + \frac{1}{1 + x} = \frac{2}{1 - x^2}$$

Operations on Generating Functions

Right Shifting

if

$$\langle f_0, f_1, f_2, f_3, \dots \rangle \longleftrightarrow F(x),$$

then

$$\langle \underbrace{0, 0, \dots, 0}_{k\text{-zeros}}, f_0, f_1, f_2, f_3, \dots \rangle \longleftrightarrow x^k F(x).$$

Operations on Generating Functions

Right Shifting

if

$$\langle f_0, f_1, f_2, f_3, \dots \rangle \longleftrightarrow F(x),$$

then

$$\langle \underbrace{0, 0, \dots, 0}_{k\text{-zeros}}, f_0, f_1, f_2, f_3, \dots \rangle \longleftrightarrow x^k F(x).$$

Derivative

if

$$\langle f_0, f_1, f_2, f_3, \dots \rangle \longleftrightarrow F(x)$$

then

$$\langle f_1, 2f_2, 3f_3, 4f_4, \dots \rangle \longleftrightarrow \frac{d}{dx} F(x).$$

Operations on Generating Functions

Example

- $\langle 1, 1, 1, 1, \dots \rangle \longleftrightarrow \frac{1}{1-x}$
- $\langle 1, 2, 3, 4, \dots \rangle \longleftrightarrow \frac{d}{dx} \frac{1}{1-x} = \frac{1}{(1-x)^2}$
- $\langle 0, 1, 2, 3, \dots \rangle \longleftrightarrow x \cdot \frac{1}{(1-x)^2} = \frac{x}{(1-x)^2}$
- $\langle 1, 4, 9, 16, \dots \rangle \longleftrightarrow \frac{d}{dx} \frac{x}{(1-x)^2} = \frac{1+x}{(1-x)^3}$
- $\langle 0, 1, 4, 9, \dots \rangle \longleftrightarrow x \cdot \frac{1+x}{(1-x)^3} = \frac{x(1+x)}{(1-x)^3}$

Operations on Generating Functions

Product

if

$$\langle a_0, a_1, a_2, a_3, \dots \rangle \longleftrightarrow A(x)$$

and

$$\langle b_0, b_1, b_2, b_3, \dots \rangle \longleftrightarrow B(x),$$

then

$$\langle c_0, c_1, c_2, c_3, \dots \rangle \longleftrightarrow A(x).B(x),$$

where

$$c_n = a_0 b_n + a_1 b_{n-1} + a_2 b_{n-2} + \dots + a_n b_0.$$

Note that $C(x) = A(x).B(x) = \sum_{n=0}^{\infty} c_n x^n$

The Fibonacci Sequence

Definition

The Fibonacci sequence is defined by the following recurrence equation:

$$f_n = \begin{cases} 0 & \text{if } n = 0, \\ 1 & \text{if } n = 1, \\ f_{n-1} + f_{n-2} & \text{if } n \geq 2. \end{cases}$$

The generating function for this sequence can be considered as follows:

$$F(x) = f_0 + f_1x + f_2x^2 + f_3x^3 + \cdots = \sum_{n=0}^{\infty} f_n x^n$$

The Fibonacci Sequence

First approach: By expanding the recurrence equation, the following sequence is obtained:

$$\langle 0, 1, f_1 + f_0, f_2 + f_1, f_3 + f_2, \dots \rangle$$

One can break this sequence into a sum of three sequences as follows:

$$\begin{array}{rcl}
 \langle 0, & 1, & 0, & 0, & 0, & \dots \rangle \longleftrightarrow x \\
 + \langle 0, & f_0, & f_1, & f_2, & f_3, & \dots \rangle \longleftrightarrow xF(x) \\
 + \langle 0, & 0, & f_0, & f_1, & f_2, & \dots \rangle \longleftrightarrow x^2F(x) \\
 \hline
 \langle 0, & 1 + f_0, & f_1 + f_0, & f_2 + f_1, & f_3 + f_2, & \dots \rangle \longleftrightarrow x + xF(x) + x^2F(x)
 \end{array}$$

$$\implies F(x) = x + xF(x) + x^2F(x) \implies F(x) = \frac{x}{1 - x - x^2}$$

The Fibonacci Sequence

Second approach:

$$f_n = f_{n-1} + f_{n-2}$$

$$\Downarrow$$

$$x^n f_n = x^n f_{n-1} + x^n f_{n-2}$$

$$\Downarrow$$

$$\sum_{n=2}^{\infty} x^n f_n = \sum_{n=2}^{\infty} x^n f_{n-1} + \sum_{n=2}^{\infty} x^n f_{n-2}$$

$$\Downarrow$$

$$\sum_{n=0}^{\infty} x^n f_n - x = x \left[\sum_{n=0}^{\infty} x^n f_n \right] + x^2 \left[\sum_{n=0}^{\infty} x^n f_n \right]$$

$$\Downarrow$$

$$F(x) - x = xF(x) + x^2 F(x)$$

$$\Downarrow$$

$$F(x) = \frac{x}{1 - x - x^2}$$

The Fibonacci Sequence

Now, we should expand $F(x)$ to extract coefficients...

$$1 - x - x^2 = (1 - \alpha_1 x)(1 - \alpha_2 x) \implies \alpha_1 = \frac{1}{2}(1 + \sqrt{5}), \alpha_2 = \frac{1}{2}(1 - \sqrt{5})$$

Next we do as follows:

$$\frac{x}{1 - x - x^2} = \frac{A_1}{1 - \alpha_1 x} + \frac{A_2}{1 - \alpha_2 x} \implies A_1 = \frac{1}{\sqrt{5}}, A_2 = -\frac{1}{\sqrt{5}}$$

The Fibonacci Sequence

Now, we should expand $F(x)$ to extract coefficients...

$$1 - x - x^2 = (1 - \alpha_1 x)(1 - \alpha_2 x) \implies \alpha_1 = \frac{1}{2}(1 + \sqrt{5}), \alpha_2 = \frac{1}{2}(1 - \sqrt{5})$$

Next we do as follows:

$$\frac{x}{1 - x - x^2} = \frac{A_1}{1 - \alpha_1 x} + \frac{A_2}{1 - \alpha_2 x} \implies A_1 = \frac{1}{\sqrt{5}}, A_2 = -\frac{1}{\sqrt{5}}$$

Now we have:

$$\begin{aligned} F(x) &= \frac{1}{\sqrt{5}} \left(\frac{1}{1 - \alpha_1 x} - \frac{1}{1 - \alpha_2 x} \right) \\ &= \frac{1}{\sqrt{5}} \left((1 + \alpha_1 x + \alpha_1^2 x^2 + \dots) - (1 + \alpha_2 x + \alpha_2^2 x^2 + \dots) \right) \end{aligned}$$

so

$$f_n = \frac{\alpha_1^n - \alpha_2^n}{\sqrt{5}} = \frac{1}{\sqrt{5}} \left(\left(\frac{1 + \sqrt{5}}{2} \right)^n - \left(\frac{1 - \sqrt{5}}{2} \right)^n \right).$$

Exercise

1. Suppose that $S_n = \{1, 2, 3, \dots, n\}$. An involution over the set S_n is a permutation $\pi : S_n \mapsto S_n$ of order at most 2 (i.e. $1 \leq \forall i \leq n, \pi^2(i) = i$). Derive a recurrence equation to count the number of involution for a set of size n and then try to solve it using Generating Functions.
2. Try to obtain a closed form for the following recurrence equation:

$$p_n = \begin{cases} 1 & \text{if } n \leq 2, \\ 2 & \text{if } n = 3, \\ p_{n-1} + (n-1)p_{n-2} - p_{n-3} + p_{n-4} & \text{if } n \geq 4. \end{cases}$$

Fast Multiplication

Suppose that x and y are integers. The size of x is assumed to be n , where n is the number of bits required to represent x ($n = \lceil \log_2(x) \rceil$). Computing $z = x \cdot y$:

Fast Multiplication

Suppose that x and y are integers. The size of x is assumed to be n , where n is the number of bits required to represent x ($n = \lceil \log_2(x) \rceil$). Computing $z = x \cdot y$:

- Direct approach: has $O(n^2)$ time complexity.

Fast Multiplication

Suppose that x and y are integers. The size of x is assumed to be n , where n is the number of bits required to represent x ($n = \lceil \log_2(x) \rceil$). Computing $z = x \cdot y$:

- Direct approach: has $O(n^2)$ time complexity.
- Consider the 1-valued bits:
 - Best case: $O(n)$ time complexity.
 - Worse case: $O(n^2)$ time complexity.

Fast Multiplication

Suppose that x and y are integers. The size of x is assumed to be n , where n is the number of bits required to represent x ($n = \lceil \log_2(x) \rceil$). Computing $z = x.y$:

- Direct approach: has $O(n^2)$ time complexity.
- Consider the 1-valued bits:
 - Best case: $O(n)$ time complexity.
 - Worse case: $O(n^2)$ time complexity.
- Divide and Conquer:

$$x \begin{array}{|c|c|} \hline \overset{n/2}{a} & \overset{n/2}{b} \\ \hline \end{array} \qquad y \begin{array}{|c|c|} \hline \overset{n/2}{c} & \overset{n/2}{d} \\ \hline \end{array}$$

Since $x = a.2^{n/2} + b$ and $y = c.2^{n/2} + d$, so

$$z = x.y = a.c.2^n + (a.d + b.c).2^{n/2} + b.d$$

In this case we have $T(n) = 4T(n/2) + O(n) = O(n^2)$! **How we can reduce the time complexity?**

Fast Multiplication

In order to improve the algorithm and reduce the time complexity, we use the following trick:

- $w_1 = a + b$
- $w_2 = c + d$
- $u = w_1 \cdot w_2 = a \cdot c + a \cdot d + b \cdot c + b \cdot d$
- $v = a \cdot c$
- $w = b \cdot d$

Now, we have:

$$z = x \cdot y = v \cdot 2^n + (u - v - w) \cdot 2^{n/2} + w$$

and so

$$T(n) = 3T(n/2) + O(n) = \Theta(n^{\log_2(3)}).$$

Fast Multiplication

```
1  Fast_Multiplication(x, y, n){
2      if(n > 1){
3          a ← MSB(x);
4          b ← LSB(x);
5          c ← MSB(y);
6          d ← LSB(y);
7          w1 ← ADD(a, b, n/2);
8          w2 ← ADD(c, d, n/2);
9          u ← Fast_Multiplication(w1, w2, n/2);
10         v ← Fast_Multiplication(a, c, n/2);
11         w ← Fast_Multiplication(b, d, n/2);
12         res ← Shift(v, n);
13         res ← Add(res, Shift(SUB(SUB(u, v, n/2), w, n/2), n/2), 2n);
14         res ← Add(res, w, 2n);
15         return(res);
16     }
17 }
```

Exercises

1. Try to solve the Fast Multiplication by dividing each number into three parts and analyze it.